

Centre de Physique Théorique* - CNRS - Luminy, Case 907
F-13288 Marseille Cedex 9 - France

Critical Properties of Stochastic Cellular Automata

Stam NICOLIS[†]

Abstract

We study the effect of mixing two rules on the dynamics of one-dimensional cellular automata by large scale numerical simulations. We calculate the decay of the magnetization for the Domany-Kinzel automaton (XOR/AND mixing) to its equilibrium value in the three phases. This requires system sizes in excess of 1 million sites. We also find severe finite size effects near the new critical points recently proposed on the basis of transfer matrix arguments.

Key-Words : Cellular Automata, Directed Percolation, Large Scale Simulations

Number of figures : 4

May 1994
CPT-94/P.3037
cond-mat/9406005

anonymous ftp or gopher: cpt.univ-mrs.fr
World Wide Web: <http://cpt.univ-mrs.fr> *or*
gopher: [//cpt.univ-mrs.fr/11/preprints](http://cpt.univ-mrs.fr/11/preprints)

*Unité Propre de Recherche 7061

[†]Centre de Physique Théorique, CNRS, Luminy, Case 907, F-13288 Marseille cedex 9, France
and PHYMAT, Université de Toulon et du Var, 83957 La Garde cedex, France

Critical Properties of Stochastic Cellular Automata

Stam Nicolis

*Centre de Physique Théorique, CNRS-Luminy, Case 907
F-13288 Marseille cedex 9, France*

ABSTRACT

We study the effect of mixing two rules on the dynamics of one-dimensional cellular automata by large scale numerical simulations. We calculate the decay of the magnetization for the Domany-Kinzel automaton (XOR/AND mixing) to its equilibrium value in the three phases. This requires system sizes in excess of 1 million sites. We also find severe finite size effects near the new critical points recently proposed on the basis of transfer matrix arguments.

Cellular automata have been studied in a systematic way for more than ten years [1] and constitute a deceptively simple family of dynamical systems-where space, time and the range of values of the dynamical variable have been discretized; yet still there are many open problems. Even better, one finds new features in well studied systems.

As a concrete illustration we shall study the Domany-Kinzel automaton [2], defined as follows:

$$\begin{aligned} P(\sigma_i^{t+1} = 1 | \sigma_{i-1}^t = 1, \sigma_{i+1}^t = 1) &= p_2 \\ P(\sigma_i^{t+1} = 1 | \sigma_{i-1}^t = 1, \sigma_{i+1}^t = 0) &= P(\sigma_i^{t+1} = 1 | \sigma_{i-1}^t = 0, \sigma_{i+1}^t = 1) = p_1 \\ P(0|1, 0) &= 1 - p_1; \quad P(0|1, 1) = 1 - p_2; \quad P(1|0, 0) = 0 \end{aligned} \quad (1)$$

starting from some initial configuration $\{\sigma_i^{t=0}\}_{i=1}^N$. If $p_1 = 1$ and $p_2 = 0$ one readily identifies the XOR rule, $P(1|1, 0) = P(1|0, 1) = 1$, $P(1|1, 1) = 0$, while, if $p_1 = 0$ and $p_2 = 1$ one has the AND rule, $P(1|1, 1) = 1$ all others giving zero.

This system has been the object of many investigations since its introduction in ref [2]-[3, 4, 5, 6], since many models of interest may be mapped onto it, in some region of its parameter space. Nevertheless many open questions remain, first among them being the accurate determination of the transition points and a study of its critical properties [7]. Whereas the original papers [2, 3] identified one critical point (from a mapping to an exactly solvable model), namely at $p_1 = 1, p_2 = 0.5$ and conjectured the existence of another one (but could not resolve the issue due to the small system size), recent work has not only—essentially—settled this issue (in the positive sense) [4, 5, 7], but has advanced the conjecture of the existence of novel critical points within the active and the chaotic phases [6].

In the present note we present results from large scale simulations on the decay of the magnetization in the various phases of the model. We also investigate what happens near the (critical) points $(p_1 = 0.614, p_2 = 0.925)$, and $(p_1 = 0.648, p_2 = 0.861)$, conjectured to exist on the basis of approximate renormalization group calculations [6].

First of all one finds that the system sizes studied to date (even those of ref. [5]) are too small to resolve this sort of question. As an example we display in fig. 1 the magnetization, $m(t)$ against time for system sizes up to 1Msites. Especially in the active and chaotic phases, equilibrium is reached so fast that it is impossible to measure how the magnetization decays to its equilibrium value. The breakpoint (in the frozen phase) appears to be 128ksites. Therefore all our runs were henceforth done at 128ksites and larger. In fact, even this size was too small to resolve many situations and 4 Msites was barely satisfactory (see below). On the other hand, In the frozen phase we find exponential decay with characteristic time $\tau \approx 80$, fig. 1.

In the active and chaotic phases, we find exponential decay, with a characteristic time essentially equal to that in the frozen phase for the active phase; the chaotic phase, on the other hand, has a significantly longer decay time, as might be expected.




Figure 1: Magnetization vs. time in the frozen phase—($p_1 = 0.3, p_2 = 0.7$) for several sizes—4096,65536,131072,262144,524288,1048576 from top to bottom.




Figure 2: Magnetization vs. time in the active phase—($p_1 = 0.75, p_2 = 0.87$) for 4194304 sites.




Figure 3: Magnetization vs. time just inside the chaotic phase—($p_1 = 0.76, p_2 = 0.2$) for 4194304 sites and two different initial conditions.

Regarding what happens at the points R and H_1 , we find that the values given for p_1, p_2 are almost surely misleading—the finite time effects from the transfer matrix method must be much too strong. For instance, for $L = 131072$ we find an exponential decay with definitely different decay times—cf. Fig. .

To summarize, we have studied the time dependence of the magnetization in the three phases of the Domany-Kinzel cellular automaton. Our results show that 4 Msites barely allow one to reach the asymptotic regime slowly enough to extract a relaxation time. Such large systems and long times are needed to resolve the differences (if they do exist) between the *true* points H_1 and R . On the other hand it should be noted that severe critical slowing down near these critical points is a problem at these sizes, as preliminary simulations have indicated. Even larger sizes are necessary to study the properties of higher correlation functions [8]—however these systems are definitely within reach of contemporary supercomputers and even fast workstations (memorywise there is no problem in running even 16 Msite systems on a Sparc 10 (or even an ELC) workstation and on a Cray YMP memory and time requirements fall within the “tiny” class). These large sizes and long times are necessary; since already, for much smaller sizes, we find, for instance, non-zero damage at $p_1 = 1, p_2 = 0.38$ (for $L = 65536$ and 1% initial damage) and for $p_1 = 0.74, p_2 = 0.0$, (for $L = 45056$ and 1% initial damage)—cf. ref. [7].

Acknowledgements: the code was developed and the first results were obtained on the Cray Y-MP of the Jülich Research Center; later results were obtained on the Cray Y-MP of the Mediterranean Institute of Technology. Computer time on these

Figure 4: Normalized magnetization, $M(t) = m(t) - m(\infty)$ vs. time for the point $(p_1 = 0.641, p_2 = 0.861)$ (point R) and the point $(p_1 = 0.614, p_2 = 0.925)$ (point H_1); $L = 131072$.

machines is gratefully acknowledged. I would also like to thank G. A. Kohring, R. Livi, H. J. Herrmann and D. Stauffer for stimulating discussions.

References

- [1] S. Wolfram, “Theory and Applications of Cellular Automata”, World Scientific (1986).
- [2] E. Domany and W. Kinzel, *Phys. Rev. Lett.* **53** 311 (1984)
- [3] W. Kinzel, *Ann. Israel Phys. Soc.* **5** 425 (1983), “Percolation Structures and Disorder”, eds. G. Deutscher, R. Zallen and J. Adler; *Z. Phys.* **B58** 229 (1985).
- [4] M. L. Martins, H. F. Verona de Rezende, C. Tsallis and A. C. N. de Magalhães, *Phys. Rev. Lett.* **66** 2045 (1991).
- [5] G. A. Kohring and M. Schreckenberg, *J. Physique I* **2** 2033 (1992).
- [6] F. Bagnoli, R. Bulajich, R. Livi and A. Maritan, *J. Phys. A: Math. Gen.* **25** L1071 (1992).
- [7] G. F. Zebende and T. J. P. Penna, *J. Stat. Phys.* **74** 1273 (1994).
- [8] S. Nicolis, (1994) in preparation.

